

NNRTIs was only confirmed after weeks of unsuccessful therapy by further resistance testing.

My patient has alerted all her friends in Nigeria as to the real nature of combivir and advised them to stop using it for breast enhancement. She has also told me that she believes the doctor in Nigeria who prescribed these drugs may have had this activity terminated.

Because no one I have spoken to has come across this particular misuse of antiretroviral therapy I felt it was worth highlighting to a wider audience in the hope that such practises may be addressed.

Acknowledgements

This patient has agreed to publish details of her case to help prevent recurrent misuse of this drug.

S M Young

Department of Genitourinary Medicine, King's Mill Hospital, Mansfield Road, Sutton in Ashfield, Nottinghamshire, NG17 4JL, UK;
Susan.Young@sfh-tr.nhs.uk

Sherwood Forest Hospitals NHS Trust Research and Development committee approval

doi: 10.1136/sti.2004.014415

Accepted for publication 21 December 2004

CORRECTION

doi: 10.1136/sti.2004.010181corr1

The order of the authors of the paper by Götz *et al* on page 24 of the February 2005 issue (HM Götz *et al*. A prediction rule for selective screening of Chlamydia trachomatis infection. *Sex Transm Infect* 2005;**81**:24–30) were wrong. The order should have been as follows: HM Götz, JEAM van Bergen, IK Veldhuijzen, J Broer, CJPA Hoebe, EW Steyerberg, AJJ Coenen, F de Groot, MJC Verhooren, DT van Schaik, and JH Richardus.